

THE STELLAR POPULATIONS OF M31 HALO SUBSTRUCTURE¹

ANNETTE M. N. FERGUSON^{2,3}, RACHEL A. JOHNSON^{4,5}, DANIEL C. FARIA^{4,6}, MIKE J. IRWIN⁷, RODRIGO A. IBATA⁸,
 KATHRYN V. JOHNSTON⁹, GERAINT F. LEWIS¹⁰, NIAL R. TANVIR¹¹

Accepted for publication in the Astrophysical Journal Letters

ABSTRACT

We present the first results from our survey of stellar substructure in the outskirts of M31 using the *Advanced Camera for Surveys* (ACS) on board the Hubble Space Telescope. We discuss the stellar populations associated with five prominent stellar overdensities discovered during the course of our panoramic ground-based imaging survey with the Isaac Newton Telescope Wide-Field Camera (INT WFC); a sixth pointing targets a region of ‘clean’ halo. The colour-magnitude diagrams (CMDs), which contain between $\approx 10,000$ – $90,000$ stars and reach four magnitudes below the horizontal branch, reveal clear variations in morphology between most fields, indicating that the age and/or metallicity mix of stars is not constant at large radius. This directly confirms the existence of large-scale population inhomogeneities within the halo of M31 and lends further support to the notion that M31 has formed, at least in part, through satellite accretions. We find a striking similarity between the populations of the giant stellar stream and those of another overdensity, the NE shelf, which lies north-east of the galaxy center. If these overdensities are associated with the same population, then the difference in their red clump magnitudes implies the NE shelf lies in front of the stream by several tens of kpc, in good agreement with recent orbit calculations for the stream progenitor.

Subject headings: galaxies: formation – galaxies: evolution – galaxies: structure – galaxies: halos – galaxies: individual (M31) – galaxies: stellar content

1. INTRODUCTION

The INT WFC survey of M31 has led to the discovery of significant stellar substructure in the halo and outer disk of our nearest giant neighbour (Ibata et al. 2001; Ferguson et al. 2002; McConnachie et al. 2003, 2004; Irwin et al. 2005). At the present time, the survey consists of more than 40 square degrees of V- and *i*-band imaging to a depth sufficient to resolve stars to ≈ 3 magnitudes below the tip of the red giant branch (RGB).

Few constraints currently exist on the nature and origin of the stellar substructure in M31. Within the popular Λ CDM model for galaxy formation, substructure is expected in the outer regions of galaxies as they continue to grow from the accretion and tidal disruption of satellite companions. Recent calculations suggest that mas-

sive disk galaxies, like the Milky Way and M31, could have cannibalised ~ 100 dwarf galaxy-like systems over their lifetimes (Bullock & Johnston 2004). Since these accreted satellites are likely to have experienced a range of star formation and chemical evolution histories, one expects their tidal debris to be distinct in terms of stellar populations. If this picture is correct, then the spatial substructure in M31 should be accompanied by stellar population inhomogeneities.

In order to gain insight into the origin of the M31 halo substructure, we are conducting a detailed study of the associated stellar populations using the HST/ACS. Our INT WFC survey indicated intriguing variations in the mean RGB colour across the halo, which could reflect changes in metallicity and/or age. HST is required to confirm these variations and, via detection of fainter and more evolved populations such as the red clump and horizontal branch, provide constraints on their nature. In this *Letter*, we present the first results from our HST/ACS survey which establish clear differences, as well as some similarities, between the populations in various substructures.

2. OBSERVATIONS

Observations at six locations in the outskirts of M31 were obtained during Cycle 11 with the Wide Field Camera of the ACS as part of GO#9458. Our fields sample the five most prominent stellar overdensities in the maps of Ferguson et al. (2002) as well as a region of apparently ‘clean’ halo (**Minor Axis**) situated 20 kpc along the southern minor axis (see Figure 1). The main characteristics of the stellar overdensities can be summarized as follows:

Giant Stream - a stellar stream, with a well-defined eastern edge, which can be traced in projection to ~ 70 kpc from the center of the galaxy towards the

¹ Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with program GO9458.

² Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Strasse 1, Garching bei München, D-85741 Germany

³ Present Address: Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh UK EH9 3HJ; ferguson@roe.ac.uk

⁴ European Southern Observatory, Alonso de Cordova 3107, Viña del Mar, Santiago, Chile

⁵ Department of Astrophysics, University of Oxford, Keble Road, Oxford UK OX1 3RH

⁶ Lund Observatory, Box 43, SE-221 00 Lund, Sweden

⁷ Institute of Astronomy, Madingley Road, Cambridge UK CB3 0HA

⁸ Observatoire de Strasbourg, 11, rue de l’Université, F-67000, Strasbourg, France

⁹ Astronomy Department, Wesleyan University, Middletown, CT 06459, USA

¹⁰ Institute of Astronomy, School of Physics, A29, University of Sydney, NSW 2006, Australia

¹¹ Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield UK AL10 9AB




FIG. 1.— A map of the surface density of blue RGB stars in a $40^\circ \times 40^\circ$ area around M31 as mapped with the INT WFC. Overlaid are the locations of the HST/ACS pointings presented here (red) and the deep HST/ACS halo pointing of Brown et al. (2003) (cyan).

south-east (Ibata et al. 2001; McConnachie et al. 2003). Radial velocity data have ruled out a simple connection to the dwarf elliptical companions M32 and NGC 205 (Ibata et al. 2004; Font et al. 2004);

NE Shelf - a large overdensity north-east of the galaxy centre which also exhibits a sharp outer boundary;

NGC 205 Loop - a small (~ 15 kpc) loop of stars which appears, in projection, to emanate from NGC 205. While the geometry is very suggestive of a connection between the two, extant stellar kinematical data still leave open other possibilities (McConnachie et al. 2004);

G1 Clump - a stellar overdensity which sits 30 kpc along the south-western major axis, in close proximity to the anomolous globular cluster G1. Although initially a connection between the two was suspected, radial velocity data have now ruled this out (Reitzel, Guhathakurta, & Rich 2004; Ferguson et al. 2004);

Northern Spur - first hinted at by Waltherbos & Kennicutt (1988), this is an extended plume of stars which sits just above the north-eastern major axis.

Each field was observed for one orbit in the F606W filter (broad V) and two orbits in the F814W filter (I), all within a single visit. Integer pixel dithers were executed between multiple sub-exposures to facilitate warm pixel

and cosmic-ray rejection. Given the low stellar density in the outer halo, two separate pointings were obtained in each of the three outermost fields (**Minor Axis**, **NE Shelf** and **Giant Stream**) in order to ensure sufficient statistics on the upper RGB and horizontal branch.

Images were first processed through the ACS pipeline and those in a given passband were combined using the Multidrizzle task (Koekemoer, Fruchter, Hook, & Hack 2002) within PyRAF. We used the default mode of Multidrizzle which calculates the offsets between dithered exposures using the information in the image header world coordinate system. The final drizzle was conducted using the Lanczos3 kernel with *pixfrac* and *scale* set to unity.

Photometry was subsequently obtained using the IRAF implementation of DAOPHOT (Stetson 1987). The stellar density in our fields is low enough to allow precise results from aperture photometry alone. PSF-fitting photometry was also carried out to compare with the aperture photometry and to measure how accurately individual sources could be modelled by a stellar PSF. For this purpose, a spatially non-varying PSF was built using ~ 150 bright stars in each of the nine ACS pointings. Sources were retained in our final photometry list if their magnitude errors, sharpnesses and chi values all lay within 3-sigma of the average value at their magnitude. Aperture corrections were derived from several tens of bright stars in each combined frame and our photometry was placed on the STMAG system using the latest zero-point values (Sirianni et al. 2005). Inspection of the Schlegel, Finkbeiner, & Davis (1998) reddening map indicates values of $E(B-V) = 0.06 - 0.08$ (with uncertainty 16%) in the far outskirts of M31; we adopt a constant value of $E(B-V) = 0.07$ for all fields and values of the selective extinction coefficient in the F606W and F814W passbands from Sirianni et al. (2005). The final CMDs contain between 10,000 stars (the **Minor Axis** field) and 90,000 stars (the **Northern Spur** field) and extend to four magnitudes below the horizontal branch (see Figure 2). Artificial star tests indicate 75% completeness to $F814W_0 = 28.5$ in fields of average stellar density. Details of our analysis will be presented in a forthcoming paper.

3. RESULTS

3.1. The CMD Morphologies

Figure 2 shows Hess diagrams for the six halo fields targetted in our survey. In all cases, a broad RGB and a prominent red clump are evident, consistent with the findings of earlier HST studies of the M31 halo (e.g. Holland, Fahlman, & Richer (1996); Bellazzini et al. (2003)). However, on closer inspection, noticeable differences can be seen in the morphology of many of the CMDs. For example, the **Giant Stream** and **NE Shelf** exhibit rather prominent RGB bumps ($F814W_0 \approx 26.1$). Note that the lower stellar density of the **Minor Axis** CMD makes it difficult to assess whether the RGB bump is similarly strong in this field. The RGB bump arises due to an evolutionary pause on the first ascent RGB when the H-burning shell passes through the discontinuity left by the deepest penetration of the convective envelope (Thomas 1967; Iben 1968). It is a short-lived phase and, although commonly seen in single-age populations, has only been recently identified in composite populations




FIG. 2.— Hess diagrams of six halo fields in M31 shown with a square root stretch. **G1 Clump**, **NGC205 Loop**, **Northern Spur**, **Giant Stream** and **NE Shelf** are regions of stellar substructure within the M31 halo, while **Minor Axis** was chosen to represent clean halo. Clear differences are apparent between most CMDs, indicating genuine stellar population variations within the halo substructure.

(e.g. Monaco, Ferraro, Bellazzini, & Pancino (2002)). The clarity of this feature in the shelf and stream CMDs suggests these regions contain a dominant stellar population with a small to moderate range of age and metallicity, perhaps the result of one or more bursts. Additionally, the location of the RGB bump at approximately 0.4 magnitudes fainter than the red clump in these fields supports the inference from the RGB colour that this is a relatively metal-rich population (e.g. Alves & Sarajedini (1999)).

The **G1 Clump** is distinct in showing clear evidence for a young population. It exhibits a well-populated upper main sequence (blue plume), with some stars as bright as $F814W_0 \approx 24.5$. Comparison to the Padova isochrones (Girardi 2004) indicates the brightest stars are consistent with a population of age of $\sim 2.5 \times 10^8$ years (Faria et al. 2005). On the other side of the major axis, and at a slightly smaller radius, the **Northern Spur** shows no evidence for such a young component. Instead, this field shows a low-amplitude RGB bump at $F814W_0 \sim 26$ and another bump at ~ 24.5 . Given the moderate metallicity inferred from the RGB colour, the models of Alves & Sarajedini (1999) indicate that the brighter bump is more likely to be an asymptotic giant branch (AGB) bump as opposed to a secondary RGB

bump (such features are also marginally present in the **G1 Clump** CMD). The **NGC 205 Loop** is similar to the **Giant Stream**, **NE Shelf** and **Minor Axis** fields in having an extended blue horizontal branch, but it differs in having neither an RGB/AGB bump nor a blue plume. Finally, all three CMDs in the top panel of Figure 2 show tantalizing evidence of a bright main-sequence turn off at $F814W_0 \approx 27.5$, roughly matched by a population of $\sim 2.5 \times 10^9$ years with $[Fe/H] = -0.7$ (Girardi 2004). As the **Giant Stream** and **NE Shelf** fields contain more stars than either the **G1 Clump** or **NGC 205 Loop**, the visibility of this putative turnoff is not simply dependent on total stellar density.

Although the CMD morphologies often differ, the mean colour of the (broad) RGB remains surprisingly constant. When measured in the region $27 < F814W_0 < 26$, the mean RGB colour varies by less than 0.05 magnitudes from region to region. This is comparable to the difference expected in colour between $[Fe/H] = -0.4$ and -0.7 stellar isochrones for a population of 14 Gyr (Girardi 2004).

3.2. Using Stellar Populations to Trace the Giant Stream

The striking resemblance between the **Giant Stream** and **NE Shelf** CMDs motivates further exploration of the relationship between these features. Figure 3 shows the $F814W_0$ luminosity function (LF) of these populations in the vicinity of the red clump and RGB bump. We fit the LF in the red clump region with the sum of a first order polynomial and a gaussian, selecting stars with the same range of de-reddened magnitude and colour in each case. We find the peak magnitude of the red clump to differ slightly between the fields, being $F814W_0 = 25.67 \pm 0.01$ for the stream and 25.53 ± 0.01 for the shelf.

The magnitude and colour of the red clump are known to vary as a function of age and metallicity (e.g. Girardi & Salaris (2001)) but the overall similarity of the CMD morphologies suggests that population-driven differences between the stream and shelf regions are small. This is further confirmed by the nearly identical luminosity function slopes observed. In this case, a more likely cause of the offset in red clump magnitude is that the regions are located at different line-of-sight distances. The red clump magnitude difference of $\Delta(F814W_0)_{\text{STREAM-SHELF}} = 0.14 \pm 0.01$ implies that the **Giant Stream** region lies beyond the **NE Shelf** by a factor of 1.07 ± 0.01 . McConnachie et al. (2003) have measured the line-of-sight distance to the stream as a function of position along it. We have interpolated their measurements to the location of our **Giant Stream** ACS field and find a line-of-sight distance of 830 ± 20 kpc. Combining this value with the differential distance modulus inferred from the red clump magnitudes yields a distance to the shelf of 776 ± 20 kpc.

4. DISCUSSION

Our HST/ACS survey of M31 substructure has revealed significant variations in CMD morphology at large radius, indicating that the age and/or metallicity of stars in these parts is not constant. This is the first direct evidence for large-scale population inhomogeneities in the outskirts of M31. Prior HST studies, which have inferred a uniform population, have generally been based on random pointings within the inner halo and have missed the main stellar overdensities (e.g. Bellazzini et al. (2003)).

It is tempting to interpret the observed population variations as evidence for multiple satellite accretions within the M31 halo, each having a distinct star formation and chemical evolution history. One puzzle, however, is the rather uniform mean RGB colour (though broad distribution) observed from field-to-field. If interpreted purely in terms of metallicity, the mean RGB colour suggests a variation of $\lesssim 0.3$ dex throughout the substructure. This could suggest that much of the halo debris comes from the accretion of a single system – one which has undergone little chemical evolution during several orbits around M31 but experienced at least one burst of star formation. On the other hand, age and metallicity may be varying in tandem throughout these regions, conspiring to produce a remarkably constant mean RGB colour (for example, as seen in the Carina dwarf spheroidal galaxy (e.g. Rizzi et al. (2003))). We will evaluate these scenarios in more detail in subsequent papers where we will model the CMDs in terms of star formation and chemical evolution histories.

Previous studies of the M31 inner halo have demonstrated that ~ 25 -50% of the population is old and




FIG. 3.— $F814W_0$ LFs in the vicinity of the red clump and RGB bump for the **Giant Stream** (left) and **NE Shelf** (right). Overlaid (dashed line) are gaussian fits to the red clump.

metal poor (e.g. Holland, Fahlman, & Richer (1996); Brown et al. (2003)). The presence of a blue horizontal branch in four of the six CMDs in Figure 2 suggests this smooth halo population is also present at larger radii. The **Northern Spur** and **G1 Clump** CMDs, which probe the most extreme radii in our sample, do not, however, show this feature. Assuming our **Minor Axis** field is representative of the smooth halo component in M31, we can ask whether we should see this component in the outermost fields studied. We find that, for both a spherical and flattened ($b/a = 0.6$) $R^{1/4}$ distribution, the amount of smooth halo expected in the **Northern Spur** and **G1 Clump** CMDs is almost entirely negligible compared to the number of stars in the overdensity (4–10% of the detected stars). In other words, the absence of an obvious extended blue horizontal branch in these fields does not necessarily mean that the smooth halo component seen at smaller radii is not also present.

The currently-favoured orbit for the stream progenitor is highly radial and passes very close to the galaxy center before re-emerging north-east of the nucleus (Ibata et al. 2004; Font et al. 2004). Although radial velocities have not yet been published for stars in the **NE Shelf** region, the strong similarity between the stream and shelf CMDs found here appears to support this prediction. Furthermore, the calculations of Ibata et al. (2004) and Font et al. (2004) indicate that the shelf region probed by our ACS pointing should lie many tens of kpc in front of our stream pointing. Our differential distance estimate based on the red clump magnitude is $\approx 50 \pm 30$ kpc and is thus in encouraging agreement.

Brown et al. (2003) have recently obtained a very deep CMD of the M31 halo at a location 11 kpc along the southern minor axis (see Figure 1). The data reveal the surprising discovery of a significant (30% by mass) intermediate-age (6–8 Gyr) metal-rich population – a result somewhat at odds with the traditional picture of stellar halos as old and metal-poor. The results presented in this *Letter* suggest caution in interpreting this finding, as well as all others that are based on observations of small single fields. The stellar populations in the outskirts of M31 exhibit large-scale variations and hence inferences about halo formation drawn from such pointings may not be representative of the entire halo. This is especially true in the case of Brown’s field, which lies in close proximity to two regions which are strongly

contaminated by stream debris.

Leo Girardi, Marco Sirianni and Tom Brown are thanked for providing results ahead of publication. AMNF acknowledges support from a Marie Curie Fellowship of the European Community under contract num-

ber HPMF-CT-2002-01758 and the ESO Santiago visitor program. Support for program GO9458 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

REFERENCES

- Alves, D. R. & Sarajedini, A. 1999, *ApJ*, 511, 225
 Bellazzini, M., Cacciari, C., Federici, L., Fusi Pecci, F., & Rich, M. 2003, *A&A*, 405, 867
 Brown, T. M., Ferguson, H. C., Smith, E., Kimble, R. A., Sweigart, A. V., Renzini, A., Rich, R. M., & VandenBerg, D. A. 2003, *ApJ*, 592, L17
 Bullock, J. S. & Johnston, K. V. 2004, in proceedings of *Satellites and Tidal Streams*, eds. F. Prada, D. Martinez-Delgado, T. Mahoney, in press (astro-ph/0401625)
 Faria, D. et al. 2005, in preparation
 Ferguson, A. M. N., Irwin, M. J., Ibata, R. A., Lewis, G. F. & Tanvir, N. R. 2002, *AJ*, 124, 1452
 Ferguson, A., Chapman, S., Ibata, R., Irwin, M., Lewis, G., & McConnachie, A. 2004, *ArXiv Astrophysics e-prints*, astro-ph/0408058
 Font, A., Johnston, K. V., Guhathakurta, P., Majewski, S. R., & Rich, R. M. 2004, *ArXiv Astrophysics e-prints*, astro-ph/0406146
 Girardi, L. 2004, private communication
 Girardi, L. & Salaris, M. 2001, *MNRAS*, 323, 109
 Holland, S., Fahlman, G. G., & Richer, H. B. 1996, *AJ*, 112, 1035
 Iben, I. 1968, *Nature*, 220, 143
 Ibata, R., Irwin, M., Lewis, G., Ferguson, A. M. N., & Tanvir, N. 2001, *Nature*, 412, 49
 Ibata, R., Chapman, S., Ferguson, A. M. N., Irwin, M., Lewis, G., & McConnachie, A. 2004, *MNRAS*, 351, 117
 Irwin, M. J., Ferguson, A. M. N., Ibata, R. A., Lewis, G. F., Tanvir, N. R. & McConnachie, A. W. 2005, in preparation
 Koekemoer, A. M., Fruchter, A. S., Hook, R. N., & Hack, W. 2002, in *The 2002 HST Calibration Workshop*, eds. S. Arribas, A. Koekemoer, and B. Whitmore (STScI: Baltimore), p.339
 McConnachie, A. W., Irwin, M. J., Ibata, R. A., Ferguson, A. M. N., Lewis, G. F., & Tanvir, N. 2003, *MNRAS*, 343, 1335
 McConnachie, A. W., Irwin, M. J., Lewis, G. F., Ibata, R. A., Chapman, S. C., Ferguson, A. M. N., & Tanvir, N. R. 2004, *MNRAS*, 351, L94
 Monaco, L., Ferraro, F. R., Bellazzini, M., & Pancino, E. 2002, *ApJ*, 578, L47
 Reitzel, D. B., Guhathakurta, P., & Rich, R. M. 2004, *AJ*, 127, 2133
 Rizzi, L., Held, E. V., Bertelli, G., & Saviane, I. 2003, *ApJ*, 589, L85
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Sirianni, M., Jee, M.J., Benitez, N., Blakeslee, J.P., Martel, A.R., Clampin, M., de Marchi, G., Ford, H.C., Gilliland, R., Hartig, G.F., Illingworth, G.D., Mack, J., McCann, W.J., & Meurer, G. 2005, *PASP*, submitted
 Stetson, P. B. 1987, *PASP*, 99, 191
 Thomas, H.-C. 1967, *Zeitschrift fur Astrophysics*, 67, 420
 Walterbos, R. A. M. & Kennicutt, R. C. 1988, *A&A*, 198, 61